

Key Advances in Cellular Hypoxia Discovery

An Infographic Timeline

Pasteur Effect

1863

Louis Pasteur observed that oxygen inhibits fermentation in yeast, indirectly introducing the idea of oxygen's effect on cellular metabolism.

Significance: First connection between oxygen levels and cellular metabolic pathways.

Otto Warburg identified the metabolic shift in cancer cells, favoring glycolysis even in the presence of oxygen (aerobic glycolysis).

Significance: Laid groundwork for understanding cellular metabolism under altered oxygen conditions.



Warburg Hypothesis

Hypoxia and Erythropoiesis

1953

Leon Jacobson and colleagues demonstrated hypoxia-induced erythropoiesis, the process of red blood cell production.

Significance: Highlighted hypoxia as a physiological regulator.

Eugene Goldwasser and colleagues purified erythropoietin (EPO), a hormone promoting red blood cell production.

Significance: Identified EPO as a hypoxia-induced hormone, linking cellular signaling to oxygen sensing.




EPO Discovery

Hypoxia Chamber

1975


Jim Billups and Barry Rothenberg develop the first Modular Incubator Chamber for controlling low oxygen cell culture environments.

Significance: Enabled low-cost hypoxia tissue culture research.



Gregg Semenza and Guang Wang published findings on hypoxia-inducible factor 1 (HIF-1).

Significance: Established HIF-1 as a key regulator of cellular response to hypoxia, controlling gene expression in low-oxygen conditions.



Discovery of HIF-1

Metabolic Reprogramming

1998


Semenza and colleagues uncovered the role of hypoxia in reprogramming cellular metabolism via HIF-1, promoting glycolysis and lactate production.

Significance: Provided insights into hypoxia's role in homeostasis, cancer and other diseases.



William Kaelin Jr. and colleagues characterized oxygen-dependent hydroxylation of HIF-1 and subsequent ubiquitination and proteasomal degradation.

Significance: Clarified the role of oxygen in regulating HIF-1 function.



HIF-1 Regulation

Mitochondrial Role in Hypoxia

2002


Navdeep Chandel and colleagues revealed that mitochondria play a central role in hypoxic signaling by producing reactive oxygen species (ROS) under low oxygen.

Significance: Connected mitochondrial function with cellular adaptation to hypoxia.



Vascular endothelial growth factor (VEGF) was identified as a hypoxia-induced mediator of angiogenesis by Napoleone Ferrara.

Significance: Demonstrated the role of hypoxia in vascular remodeling and its relevance to diseases like cancer.



Role of VEGF

Nobel Prize

2019

William Kaelin Jr., Sir Peter Ratcliffe, and Gregg Semenza received the Nobel Prize for their groundbreaking discoveries related to oxygen-sensing mechanisms. All three laureates acknowledged using the Modular Incubator Chamber to create hypoxic environments in their research.

Significance: Recognized decades of work revealing how cells adapt to oxygen availability, cementing the field's importance.



FDA approved belzutifan, a HIF-2 inhibitor, for treating cancers like renal cell carcinoma.

Significance: Marked the transition of hypoxia research into therapeutic applications.




HIF-2 Inhibitor Approved

Recent Advances

2020+

Many researchers continue advancing our understanding of hypoxic mechanisms, particularly in immune regulation, stem cell biology, infectious disease, and tissue regeneration, leading to improved cell culture modeling, optimized therapeutic cell cultures, and targeting hypoxia in disease.

Significance: Results have established the wide-reaching influence of hypoxia on cellular processes, with implications for therapies in ischemia, cancer, and beyond.



Embrient Inc

Formerly known as **billups-rothenberg inc**

Empowering Reproducibility in Cell Culture

embrient.com